

Analysis of Patients Receiving Treatment for Inflammatory Breast Disease at Surgery Department of Tertiary Care Units

Muhammad Rashid¹, Muhammad Nasir Shahbaz², Amna Akram³, Awais Anwar⁴, Muhammad Umar⁵, Muhammad Shazib Ali⁶, Asfand Yar Mujahid⁷

1. Senior Registrar Surgery Sir Gangaram Ram Hospital Lahore Pakistan
2. Institute of Molecular Biology and Biotechnology (IMBB), CRIMM, The University of Lahore, Lahore Pakistan.
3. Assistant Professor Department of Biochemistry Sahara Medical College Narowal Pakistan.
4. Professor. Department of physiology Sahara Medical College Narowal Pakistan.
5. Associate Professor Department of Anatomy Nawaz Sharif Medical College, Gujrat, Pakistan.
6. Demonstrator Department of Biochemistry Rashid Latif khan university Ferozepur Road Lahore Pakistan.
7. Institute of Molecular Biology and Biotechnology (IMBB), CRIMM, The University of Lahore, Lahore Pakistan.



*Corresponding Author: Muhammad Rashid, Email: mohammad_rasheed@hotmail.com, Cell: +923324511067

ABSTRACT

Background: In the Surgery Department of Tertiary Care Units, an analysis of patients receiving treatment for inflammatory breast disease presents a critical assessment of disease patterns and therapeutic outcomes. This evaluation is pivotal in understanding the prevalence and efficacy of treatment modalities in addressing complex breast conditions within specialized care settings.

Aims and Objectives: To assess the effectiveness of the care given to mastitis and breast abscess patients admitted to the surgical department of the Sir Gangaram Ram Hospital Lahore.

Methodology: Prospective Analytical Cross-sectional Study. Current study was conducted in surgical department of the Sir Gangaram Ram Hospital Lahore from March 2023 to December 2023. The patients those visited and admitted with a diagnosis of breast abscess were included regardless of age, size of the abscess and co-morbidities. The patient was informed of the available treatment options, and their priority was taken into account for the chosen course of action. Total 100 female patients were divided in two different groups. In group-A 60 patients were operated with local anesthesia and their abscess size was larger than 2.5 cm in diameter. While in group-B, 40 patients were conservatively managed and either ultrasound guided aspiration or no intervention is used. Either the size of the abscess was less than 2.5 centimeters in these patients, or there were signs of inflammation without any fluctuation. Whereas in case of conservative treatment hot fomentation and antibiotics were recommended, some patients who received surgery or minimally invasive treatment also received prescriptions for antibiotics. The antibiotics used were a combination of a beta-lactamase inhibitor, clavulanic acid, and a broad-spectrum penicillin, namely amoxicillin, which could be used as an adjuvant or sole therapy. If there was no need for surgery to treat the abscess, the treatment was considered successful. Bio- statistical program (SPSS) version 27.0 were used for the raw data description.

Results: Total 100 female patients were considered and their aged from 20-40 years, lactating women were 42 while 58 women non-lactating. Group A consisted of 48 surgical patients who underwent incision and drainage, while Group B consisted of 52 patients who were treated conservatively. Ischemia, skin excoriation, wound sepsis, and postoperative septicemia were among the complications that were encountered during the postoperative period. Breast deformity, sinus formation, non-healing or closure, and organized abscess were diagnosed during the follow-up. There was no any mortality.

Conclusion: Surgical drainage is the only option for treating breast abscesses, both large and small, as it significantly reduces morbidity.

Keywords: Breast abscesses, lactating women, inflammation, antibiotics, Ischemia, skin excoriation, wound sepsis.

dmls



Received: 02/02/2024
Revised: 12/03/2024
Accepted: 19/03/2024
Published: 31/03/2024

© The Author(s) 2024. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third-party material in this article are included in the article's Creative Commons license unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you must obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/public-domain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

INTRODUCTION

A localized collection of pus (inflammatory exudate) in the breast tissue is called a breast abscess. Breast abscesses can occur as the initial symptom of a breast infection, but they most frequently occur when mastitis or cellulitis is unresponsive to antibiotic therapy. With a documented incidence of 0.1 percent, it is an uncommon issue for nursing mothers; among those who have had mastitis treated with antibiotics, the incidence is 3 percent. Mastitis can result in complications such as breast abscesses, which can develop from acute bacterial mastitis into a pyogenic abscess if antibiotic treatment is unsuccessful[1]. Breast abscesses are more common in non-puerperal mastitis than in puerperal mastitis, and because of their extreme discomfort and propensity to recur, they can be especially challenging to treat[2].

Historically, surgical incisions have been necessary for breast abscesses in order to drain the abscess; this procedure is typically carried out under general anesthesia and is followed by the administration of antibiotics[3]. Nevertheless, this course of treatment may cause problems with lactation and lead to unsatisfactory cosmetic outcomes[4]. More recently, it has been demonstrated that breast abscesses can be successfully treated with US-guided needle aspiration or catheter drainage. The purpose of this study was to evaluate the efficacy of minimally invasive treatments for breast abscesses by retrospectively analyzing diagnostic results, various treatment modalities, and patient outcomes treated by a multimodality breast team at our institution. In the past, incision and drainage of abscesses was thought to be the recommended course of treatment[5, 6].

This approach is more invasive than needle aspiration and often leaves scarring with structural damage and subpar cosmetic results, even though the recurrence rate is lower. While oral ant-staphylococcal antibiotics and serial aspiration have been shown to be successful, surgical excision is sometimes necessary for infected or obstructed lactiferous ducts and has been shown to reduce the recurrence rate of non-puerperal abscess and mastitis[7]. A bacterial infection is typically the cause of a breast abscess. The bacteria cause a breast infection known as mastitis when they penetrate through a break or crack in the skin of the breast or nipple. This is an inflammation that affects nursing women's breast tissue, milk glands, or ducts. An abscess is a pus-filled empty space in the breast caused by infected milk ducts[8, 9].

MATERIALS AND METHODS

This study aimed to assess the effectiveness of the care provided to patients with mastitis and breast abscesses admitted to the surgical department of Sir Gangaram Ram Hospital Lahore. The study was conducted from March

2023 to December 2023 and utilized a prospective analytical cross-sectional design.

The study included female patients diagnosed with breast abscesses, regardless of their age, abscess size, or co-morbidities. The diagnosis of inflammatory breast disease was confirmed for each patient, and the appropriateness of the treatment options in the tertiary care unit was assessed, including general health, organ function, and the absence of contraindications to therapy. Cancer patients and pregnant women were excluded from the study to eliminate confounding factors. Furthermore, patients with prior malignancies within a specific timeframe prior to diagnosis were excluded to prevent any influence on the treatment outcomes.

A total of 100 female patients were included in the study and divided into two groups. Group A consisted of 60 patients who underwent surgery with local anesthesia for abscesses larger than 2.5 cm in diameter. Group B included 40 patients who were conservatively managed with either ultrasound-guided aspiration or no intervention. The patients in Group B either had abscesses smaller than 2.5 cm or had signs of inflammation without fluctuation. Conservative treatment involved hot fomentation and antibiotics, while some patients who underwent surgery or minimally invasive treatments also received antibiotics. The antibiotics used in the study were a combination of clavulanic acid (a beta-lactamase inhibitor) and amoxicillin (a broad-spectrum penicillin), which could be used as sole therapy or an adjunct to treatment. If the abscess did not require surgery and showed resolution, the treatment was considered successful. Clinical and demographic data were recorded in MS Excel, and statistical analysis was performed using SPSS version 27.0. This study adhered to ethical standards in accordance with the Declaration of Helsinki. It was approved by the Institutional Review Board (IRB). Informed consent was obtained from all participants, ensuring they were fully aware of the nature of the study, potential risks, and available treatment options. Patient confidentiality was maintained by anonymizing all data. Participants had the right to withdraw from the study at any time without any impact on their medical treatment. All interventions and procedures followed were in line with ethical medical practices, ensuring the safety and well-being of the participants.

RESULTS

A breast abscess is a localized infection and pus buildup in the breast parenchyma. Most often, a breast abscess is a consequence of mastitis. It is the most common breast pathology seen in surgical outpatient departments. It is more common in women from lower socioeconomic backgrounds and less common in those with western lactiferous ducts, which can cause milk stasis and

infection. Most of the time, *Staphylococcus aureus* is the causing agent. The majority of non-puerperal abscesses are seen in postmenopausal women and are uncommon. It can be brought on by nipple inversion, congenital anomalies, duct ectasia, or an underlying cancer. Obesity diabetes, and smoking are risk factors. It happens when keratotic debris obstructs the lactiferous ducts, causing organisms trapped in the ducts to multiply. The most common causes of non-puerperal abscesses are anaerobes, staphylococci, and streptococcus.

Warm or cold compresses, antibiotic therapy for 10–15 days, and continuous breast emptying through breastfeeding or breast pumping every two hours or when engorged are the standard treatments for mastitis. It has been demonstrated that antibiotic therapy combined with continuous breast emptying is more effective than breast emptying alone at relieving symptoms, lowering the chance of recurrence, and lowering the risk of abscess formation. In group-A 60 patients were operated with local anesthesia and their abscess size was larger than 2.5 cm in diameter, shown in table-2.

Table-1: General information (n=100)

Demographics	Mean $\pm$ SD	P< (0.05)
Age	38.13 $\pm$ 0.01	0.01
Gender (women)	100.00 $\pm$ 0.01	0.01
Lactating women	42.01 $\pm$ 0.04	0.04
Non-Lactating women	58.01 $\pm$ 0.01	0.01
Married	78.02 $\pm$ 0.02	0.02
Unmarried	22.01 $\pm$ 0.01	0.01
Patients operated with general anesthesia	60.01 $\pm$ 0.01	0.01
Patients treated with conservative treatment	40.01 $\pm$ 0.01	0.01

Table-2: Patients operated with local anesthesia for abscess surgery (n=60)

Parameters	%(Mean $\pm$ SD)	P< (0.05)
Non-lactating abscess	18.10 $\pm$ 0.01	0.01
Lactating abscess	40.02 $\pm$ 0.01	0.01
Non-puerperal abscess	12.01 $\pm$ 0.01	0.01
puerperal abscess	48.02 $\pm$ 0.01	0.01
Simple Lactating abscess	50.01 $\pm$ 0.03	0.03
Complicated Lactating abscess	21.01 $\pm$ 0.01	0.01
Simple non Lactating abscess	20.01 $\pm$ 0.02	0.02
Complicated non Lactating abscess	10.01 $\pm$ 0.01	0.01

Table-3: Patients Treated with conservative treatment (n=40)

Parameters	%(Mean $\pm$ SD)	P< (0.05)
Non-lactating abscess	11.10 $\pm$ 0.01	0.01
Lactating abscess	30.02 $\pm$ 0.01	0.01
Non-puerperal abscess	02.01 $\pm$ 0.01	0.01
puerperal abscess	08.02 $\pm$ 0.01	0.01
Simple Lactating abscess	20.01 $\pm$ 0.01	0.01
Complicated Lactating abscess	01.01 $\pm$ 0.01	0.01
Simple non Lactating abscess	10.01 $\pm$ 0.01	0.01
Complicated non Lactating abscess	15.01 $\pm$ 0.02	0.02




Figure 1: The mean percentages with standard deviations for various abscess types. Each bar represents the mean percentage for a specific abscess category, and the error bars indicate the standard deviation. Categories where (p < 0.05) are marked with an asterisk, denoting statistical significance.




Figure 2: The mean percentages with standard deviations for various abscess categories, along with their significance levels indicated by asterisks where the p-value is less than ($p < 0.05$).

In our investigation, we found that Group-B had a higher incidence of breast deformity as a result of organized abscesses, and those who needed surgical drainage as a result of conservative treatment failure were included in the study.

Due to low immunity and co-morbid conditions, late complications did not correlate with age, but early complications were found to be strongly correlated with age. According to the study, there is a significant correlation between primary breast abscess and its recurrence and malnutrition, particularly anemia, multiparity, and poor hygiene

DISCUSSION

In present study, we found that 18% of patients in Group I had hypertrophic scars and scarring that caused contractures, and 12% of patients in Group II had breast deformities[10]. 10% of the women in Group A of our study experienced pus discharge after incision and drainage, and this finding was replicated in another study where 10% of the women experienced persistent discharge for approximately six weeks[11, 12]. In our study, 1 in 12 of the women in group I needed additional surgery. A retrospective investigation on 53 female patients with breast abscesses was carried out. Twenty-two abscesses were aspirated; five of them needed to be cut open and drained later. Thirteen patients underwent primary incision and drainage, with one patient requiring additional drainage[13, 14].

Afridi SP et al. (2013) conducted a study on the aspiration of breast abscess and found that 43.6% of patients in group II in our study required a second aspiration, while 43% of women required one after the first aspiration. Various researchers conducted a study on the diagnosis, treatment, and outcome of breast abscesses[15, 16]. One patient out of every 56 had a seroma. Another patient developed a hematoma where the drainage catheter was inserted. In our investigation, we found that group B had a higher incidence of breast

deformity as a result of organized abscesses, and those who needed surgical drainage as a result of conservative treatment failure were included in the study[17, 18, 19].

Our study found no mortality, which we attributed to early presentations, efficient antibiotic therapy, and most importantly, prompt intervention[20]. Due to low immunity and co-morbid conditions, late complications did not correlate with age, but early complications were found to be strongly correlated with age. According to the study, there is a significant correlation between primary breast abscess and its recurrence and malnutrition, particularly anemia, multiparity, and poor hygiene[21].

CONCLUSION

Even though there are benefits to not having to undergo anesthesia or endure uncomfortable dressing changes, conservative treatment for small breast abscesses such as minimally invasive ultrasound guided aspirations increase morbidity and disfigurement and may even require surgery. Both large and small breast abscesses require surgical drainage as the only viable course of treatment.

Acknowledgment: We would like to acknowledge our fellow colleagues and our departmental staff for helping and motivating us.

Authors Contributions: All authors contributed equally and sincerely in present research.

Funding: No external funding was received for the Research.

Conflict of Interest: No conflict of Interest was declared by any author during the current study.

REFERENCES

1. Enkobahry A, Sime T, Kene K, Mateos T, Dilnesa S, Zawdie B. Blood biomarkers as potential malnutrition screening alternatives among adult patients with cancer on treatment in oncology unit of Jimma tertiary hospital: A cross-sectional analysis. *BMC Nutr.* 2023;9(1):38.
2. Taylor GB, Paviour SD, MUSAAD S, Jones WO, Holland DJ. A clinicopathological review of 34 cases of inflammatory breast disease showing an association between corynebacteria infection and granulomatous mastitis. *Pathology.* 2003;35(2):109–19.

3. Jafar D, Pesnani F, Arif S, Nasir A. Self-esteem of patients receiving chemotherapy treatment in a tertiary care hospital Karachi, Pakistan. *Pak J Med Dent*. 2020;9(3):60–4.
4. Ladha K, Melo MFV, McLean DJ, Wanderer JP, Grabitz SD, Kurth T, et al. Intraoperative protective mechanical ventilation and risk of postoperative respiratory complications: hospital based registry study. *BMJ*. 2015;351:h3646.
5. Chatterjee S, Chattopadhyay A, Todi SK. Outcomes of severe sepsis among adults in a tertiary care hospital in Kolkata – a preliminary study. *J Anesth Intensive Care Med*. 2018;6(2):555684.
6. Lal S, Khan E, Malik MA. Practical challenges of managing COVID-19 positive patients in the adult intensive care unit of a tertiary care teaching hospital of Ireland. *Anaesth Pain Intensive Care*. 2020;24(3):263–71.
7. Partridge J, Harari D, Martin F, Peacock J, Bell R, Mohammed A, et al. Randomized clinical trial of comprehensive geriatric assessment and optimization in vascular surgery. *Br J Surg*. 2017;104(6):679–87.
8. Motsisim S, Ballot DE. A review of necrotising enterocolitis in very low birth weight babies in a tertiary hospital in Johannesburg. *Afr J Paediatr Surg*. 2023;20(1):59–66.
9. Garassino MC, Whisenant JG, Huang LC, Trama A, Torri V, Agustoni F, et al. COVID-19 in patients with thoracic malignancies (TERAVOLT): first results of an international, registry-based cohort study. *Lancet Oncol*. 2020;21(7):914–22.
10. Hamieh NM, Akel R, Anouti B, Traboulsi C, Makki I, Hamieh L, et al. Cancer-related pain: prevalence, severity and management in a tertiary care center in the Middle East. *Asian Pac J Cancer Prev*. 2018;19(3):769–75.
11. McGillicuddy E, Schuster K, Barre K, Suarez L, Hall M, Kaml G, et al. Non-operative management of acute cholecystitis in the elderly. *Br J Surg*. 2012;99(9):1254–61.
12. Mei AHY, Jin WLC, Hwang MKY, Meng YC, Seng LC, Yaw WH. Value of the Palliative Performance Scale in prognostication of advanced cancer patients in a tertiary care setting. *J Palliat Med*. 2013;16(8):887–93.
13. Low JA, Berman AW, Steinberg SM, Danforth DN, Lippman ME, Swain SM. Long-term follow-up for locally advanced and inflammatory breast cancer patients treated with multimodality therapy. *J Clin Oncol*. 2004;22(20):4067–74.
14. Matthews LR, Ahmed Y, Wilson KL, Griggs DD, Danner OK. Worsening severity of vitamin D deficiency is associated with increased length of stay, SICU cost, and mortality rate in surgical ICU patients. *Am J Surg*. 2012;204(1):37–43.
15. Lim B, Woodward WA, Wang X, Reuben JM, Ueno NT. Inflammatory breast cancer biology: the tumour microenvironment is key. *Nat Rev Cancer*. 2018;18(8):485–99.
16. Chaparro M, Verreth A, Lobaton T, Gravito-Soares E, Julsgaard M, Savarino E, et al. Long-term safety of in utero exposure to anti-TNF α drugs for the treatment of inflammatory bowel disease: results from the TEDDY study. *Am J Gastroenterol*. 2018;113(3):396–403.
17. Wolfrum A, Kümmler S, Theuerkauf I, Pelz E, Reinisch M. Granulomatous mastitis: a therapeutic and diagnostic challenge. *Breast Care (Basel)*. 2018;13(6):413–8.
18. He Y, Wong YWY, Ngeow AJH, Sim EY, Cherng BPZ, Arunachalam S, et al. Establishment of an isolation maternity unit in a tertiary hospital during COVID-19 pandemic. *BMC Pregnancy Childbirth*. 2022;22(1):317.
19. Senkus E, Kyriakides S, Ohno S, Penault-Llorca F, Poortmans P, Rutgers E, et al. Primary breast cancer: ESMO Clinical Practice Guidelines. *Ann Oncol*. 2015;26(Suppl 5):v8–30.
20. Kepkey BB. Barriers and facilitators of adherence to IV chemotherapy among breast cancer patients at a tertiary hospital in the Western Cape. Stellenbosch University; 2022. (*Thesis—no journal abbreviation*)
21. Moletta L, Pierobon ES, Capovilla G, Costantini M, Salvador R, Merigliano S, et al. International guidelines and recommendations for surgery during COVID-19 pandemic: systematic review. *Int J Surg*. 2020;79:180–8.

This article may be cited as: Rashid M, Shahbaz MN, Akram A, Anwar A, Umar M, Ali MS, Mujahid AY. Analysis of patients receiving treatment for inflammatory breast disease at surgery department of tertiary care units. *Dev Med Life Sci*. 2024;1(1):2–6. doi:10.69750/dmls.01.01.012.

Publisher's Note:

Developmental Medico-Life-Sciences remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

