


Prevalence and Risk Factors of Polycystic Ovary Syndrome Among Adolescent Girls in Urban and Rural Areas of Punjab, Pakistan

Sara Waheed Butt ^{1*}, Umer Farooq ², Tahreem Waheed Butt ³, Abdullah Nawaz Rai ⁴, Ammar Anjum ⁵

- 1- House Officer, Chaudhry Muhammad Akram Teaching & Research Hospital, Lahore, Pakistan
- 2- House officer, Department of Surgery, Farooq Hospital, Lahore, Pakistan
- 3- Medical Officer, Farooq Hospital, Lahore, Pakistan
- 4- House Officer, Department of Radiology, Services Hospital, Lahore, Pakistan
- 5- Medical Officer, Omar Hospital and cardiac center, Lahore, Pakistan



*Corresponding Author: Sara Waheed Butt Email: sarahwaheedbutt03@gmail.com Cell: +92 333 0497069

ABSTRACT

Background: Polycystic ovary syndrome (PCOS) is a leading endocrine disorder during adolescence, conferring reproductive, metabolic, and psychological burdens. Reliable estimates from Pakistan—particularly contrasting urban and rural settings—remain scarce.

Objective: To estimate the prevalence of PCOS and delineate its principal risk correlates among adolescent girls residing in urban and rural districts of Lahore, Pakistan.

Methods: A cross-sectional survey was undertaken between January and December 2024. Sixty girls aged 13–19 years were recruited by purposive sampling, with equal representation from urban and rural communities. Standardised questionnaires captured sociodemographic and lifestyle variables; physical and anthropometric assessments, transabdominal ultrasonography, and fasting biochemical profiles were obtained. PCOS was defined according to the Rotterdam criteria. Data were analysed with SPSS (version 25.0).

Results: The overall prevalence of PCOS was 26.7%. Urban participants demonstrated a significantly higher prevalence than their rural counterparts (36.7% vs 16.7%; $p = 0.04$). Urban residency was associated with elevated body-mass index, greater waist circumference, a higher proportion of obesity, sedentary behaviour, and frequent fast-food intake. Menstrual irregularity, acne, and hirsutism were the predominant clinical manifestations. Biochemically, girls with PCOS exhibited raised serum testosterone, fasting insulin, fasting glucose, and HOMA-IR indices (all $p < 0.05$). A positive familial history of PCOS was more common among urban respondents.

Conclusion: PCOS affects more than one in four adolescent girls in Lahore, with a markedly greater burden in urban communities driven by adiposity, lifestyle patterns, and hereditary predisposition. Early detection, focused health education, and targeted lifestyle modification programmes are essential to mitigate future cardiometabolic and reproductive sequelae.

Keywords: Polycystic, Adolescents, Urban, Rural, Obesity, Insulin



©The Author(s) 2025. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third-party material in this article are included in the article's Creative Commons licence unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you must obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Received: 31/03/2025
Revised: 28/04/2025
Accepted: 06/05/2025
Published: 07/05/2025

INTRODUCTION

Polycystic ovary syndrome (PCOS) is a common and complex endocrine disorder that is now the most common cause of menstrual irregularities, androgens and infertility in women of reproductive age everywhere [1]. PCOS usually presents clinically with irregular or absent menstrual periods, excessive hair growth, acne, obesity, insulin resistance, and multiple small cysts of the ovary. PCOS has been accepted as an adult condition for a long time, but it is now increasingly recognized as a condition that starts in adolescence. Because PCOS can lead to long term reproductive, metabolic and psychological complications, including type 2 diabetes, cardiovascular disease, infertility, depression and anxiety, it is important to identify early in life and manage [2, 3].

Across different countries and ethnic groups, the prevalence of PCOS shows considerable variation, largely due to differences in diagnostic criteria, genetic makeup, environmental exposures, and lifestyle practices. According to international estimates, PCOS affects between 6% and 20% of women and this burden is increasing in South Asian countries, including Pakistan. This upward trend has been attributed to factors including rapid urbanization, changes in the diet characterized by increased consumption of high calorie and processed foods, decreased levels of physical activity and increased incidence of obesity in children and adolescents. Despite a high prevalence of PCOS in Pakistan, most of the research has been done in adult women and very little is known about prevalence and risk factors of PCOS in adolescents, especially in the rural areas where access to healthcare and reproductive health awareness is minimal [4, 5].

Timing of intervention during adolescence is critical because it is a period of time when PCOS can be prevented or delayed through lifestyle modification, counseling, and

medical intervention when needed [6]. However, in Pakistan the diagnosis of PCOS is often delayed due to a social stigma, a lack of awareness, no screening programs and no specialized health services in rural areas. The distribution and determinants of PCOS in urban and rural adolescent populations need to be understood in order to address this gap [7, 8].

The current study was designed to estimate the prevalence of PCOS among adolescent girls in selected urban and rural regions of Punjab, Pakistan, and to investigate the key demographic, lifestyle, and clinical factors associated with its occurrence. By providing comparative data from these two distinct settings, this current study aimed to inform public health strategies, enhance early detection efforts, and ultimately improve the reproductive and metabolic health of adolescent girls in the Pakistani context [9, 10].

MATERIALS AND METHODS

This cross-sectional study was carried from January 2024 to December 2024 in Lahore, Punjab, Pakistan. The main aim of the study was to estimate the prevalence of polycystic ovary syndrome (PCOS) and its associated risk factors among adolescent girls residing in the city in both urban and rural areas. The study received the Institutional Review Board (IRB) approval. All of the participants gave written informed consent from the parents and legal guardians or verbal assent from the adolescent girls. All participants were maintained in confidentiality and anonymity and participation was voluntary and the participants could withdraw anytime without any adverse consequences.

The study population included adolescent girls, aged 13–19 years with menarche who were residing in their own area for at least 1 year. However, girls who had been

diagnosed with known endocrine disorders including hypothyroidism, hyperprolactinemia, Cushing's syndrome, or congenital adrenal hyperplasia, or were using hormonal medications or oral contraceptive pills at the time of enrollment were excluded. Purposive sampling was used to recruit an equal number of 60 girls from urban and rural localities of Lahore to have adequate representation of both the groups and the final sample size was 60 girls.

Pretested, structured questionnaire administered through face-to-face interviews by trained female researchers in private setting was used to collect data. Sociodemographic data such as age, residence, educational status, family income, and parental education were captured in the questionnaire. Menstrual cycle length, duration of flow, age at menarche and history of oligomenorrhea or amenorrhea were obtained from menstrual history. The clinical features of hyperandrogenism were determined by the presence and severity of hirsutism (modified Ferriman and Gallwey score), acne, and androgenic alopecia. Questions on physical activity level, duration of sedentary activities such as television or mobile phone use, dietary patterns, including how often fast food is consumed and sleep habits were used to evaluate lifestyle factors. PCOS, diabetes, obesity and cardiovascular diseases were also recorded in the family history.

A thorough physical examination was performed to record anthropometric measurements, including height, weight, body mass index (BMI), waist circumference, hip circumference, and waist-to-hip ratio, following standard protocols. Blood pressure was measured in a seated position using a calibrated digital sphygmomanometer. Clinical signs of insulin resistance, such as acanthosis nigricans, were noted. Transabdominal pelvic ultrasonography was conducted by an experienced sonographer using a portable

ultrasound machine to evaluate ovarian morphology, specifically the presence of polycystic ovaries characterized by the presence of 12 or more small follicles (2–9 mm in diameter) arranged peripherally around a dense stroma.

Fasting morning venous specimens were collected after a 10–12-hour fast to quantify serum total testosterone, fasting insulin, and fasting plasma glucose. Samples were processed immediately in a central, accredited laboratory, where analytes were measured with validated chemiluminescent immunoassays and enzymatic methods. Insulin resistance was estimated using the homeostatic model assessment (HOMA-IR). Polycystic ovary syndrome was diagnosed according to the Rotterdam consensus, which stipulates that at least two of three criteria—oligo- or anovulation, clinical or biochemical hyperandrogenism, and polycystic ovarian morphology on ultrasound—must be present after exclusion of alternative disorders.

Data were entered into SPSS (version 25.0). Continuous variables—including age, body-mass index, and hormone concentrations—are presented as means $\pm$ standard deviations, and categorical variables—including menstrual irregularity, hirsutism, and family history—are reported as frequencies and percentages. Urban–rural differences were assessed with independent-sample *t* tests for continuous data and χ^2 or Fisher's exact tests for categorical data. Multivariable logistic regression, adjusting for covariates identified in univariable analyses, was employed to identify independent predictors of PCOS. Two-sided *p* values < 0.05 were considered statistically significant.

RESULTS

This study included a total of 60 adolescent girls aged 13–19 years from urban and rural

areas of Lahore, with an equal distribution of 30 girls from each area. The overall prevalence of polycystic ovary syndrome (PCOS), diagnosed according to the Rotterdam criteria, was 26.7% (16 out of 60), with urban girls showing a significantly higher prevalence (36.7%, 11 out of 30) compared to rural girls (16.7%, 5 out of 30; $p = 0.04$).

Demographic and Socioeconomic Characteristics:

The mean age of participants was 16.1 ± 1.7 years, with no statistically significant difference between urban and rural groups. Educational status differed significantly between the groups, with a higher proportion of urban girls enrolled

in high school, while rural girls had higher middle school enrollment ($p = 0.04$).

Family income levels were markedly different between urban and rural participants ($p = 0.01$), with most rural families earning less than 100,000 PKR per month, whereas urban families were largely in the middle- or higher-income brackets. Parental education also showed a clear urban-rural disparity, with more urban mothers and fathers being graduates compared to their rural counterparts. Family size was larger in rural households (6.4 ± 1.3) compared to urban households (5.2 ± 1.1 ; $p = 0.001$). Most families in both groups lived in owned houses, although ownership was slightly higher among rural families.

Table 1: Demographic and Socioeconomic Characteristics of Participants

Characteristic	Urban (n=30)	Rural (n=30)	p-value
Mean age (years)	16.3 ± 1.8	15.9 ± 1.6	0.34
Education level (%)			0.04*
- Middle school	20% (n=6)	43.3% (n=13)	
- High school	80% (n=24)	56.7% (n=17)	
Monthly family income (%)			0.01*
- <100,000 PKR	26.7% (n=8)	66.7% (n=20)	
- 100,000–300,000 PKR	50% (n=15)	30% (n=9)	
- >300,000 PKR	23.3% (n=7)	3.3% (n=1)	
Mother's education (%)			0.02*
- Primary	10% (n=3)	36.7% (n=11)	
- Secondary	46.7% (n=14)	50% (n=15)	
- Graduate	43.3% (n=13)	13.3% (n=4)	
Father's education (%)			0.01*
- Secondary	30% (n=9)	60% (n=18)	
- Graduate	70% (n=21)	40% (n=12)	
Family size (mean)	5.2 ± 1.1	6.4 ± 1.3	0.001*
Housing status (%)			0.05*
- Owned house	86.7% (n=26)	96.7% (n=29)	
- Rented house	13.3% (n=4)	3.3% (n=1)	

*Statistically significant ($p < 0.05$)

Lifestyle and Anthropometric Characteristics:

Urban girls had significantly higher body mass index (BMI) and waist circumference compared to rural girls. Obesity (BMI ≥ 25 kg/m²) was more prevalent among urban participants (40%) versus rural participants (16.7%; $p = 0.03$). Sedentary lifestyle, defined

as less than 150 minutes of moderate physical activity per week, was significantly more common in urban girls (60%) than in rural girls (33.3%; $p = 0.02$). Fast-food consumption (≥ 3 times per week) was reported more frequently by urban participants (50%) compared to rural participants (23.3%; $p = 0.03$), while rural girls engaged more in physical activity.

Table 2: Lifestyle and Anthropometric Characteristics

Characteristic	Urban (n=30)	Rural (n=30)	p-value
BMI (kg/m²)	24.8 $\pm$ 3.1	22.1 $\pm$ 2.7	0.01*
Obesity (%)	40% (n=12)	16.7% (n=5)	0.03*
Waist circumference (cm)	82.4 $\pm$ 7.3	76.2 $\pm$ 6.9	0.01*
Sedentary lifestyle (%)	60% (n=18)	33.3% (n=10)	0.02*
Fast-food ≥ 3x/week (%)	50% (n=15)	23.3% (n=7)	0.03*
Physical activity >150 min/week (%)	30% (n=9)	53.3% (n=16)	0.04*

Prevalence and Clinical Features of PCOS:

PCOS was diagnosed in 26.7% of the overall study population, with a higher prevalence in urban girls. Menstrual irregularities were the most common presenting feature in both

groups, followed by acne, hirsutism, and acanthosis nigricans. Positive family history of PCOS was reported more often by urban girls (45.5%) compared to rural girls (20%), though this difference was not statistically significant.

Table 3: Prevalence and Clinical Features of PCOS

Feature	Urban PCOS (n=11)	Rural PCOS (n=5)	p-value
Oligomenorrhea (%)	63.6% (n=7)	60% (n=3)	0.86
Amenorrhea (%)	18.2% (n=2)	20% (n=1)	0.90
Hirsutism (%)	54.5% (n=6)	40% (n=2)	0.56
Acne (%)	72.7% (n=8)	60% (n=3)	0.60
Acanthosis nigricans (%)	36.4% (n=4)	20% (n=1)	0.50
Family history of PCOS (%)	45.5% (n=5)	20% (n=1)	0.20

Biochemical Profile:

Biochemical parameters demonstrated significantly elevated serum testosterone levels, fasting insulin, fasting glucose, and HOMA-IR scores in PCOS participants compared to non-PCOS participants, indicating the presence of metabolic disturbances and insulin resistance.

Table 4: Biochemical Parameters in PCOS and Non-PCOS Groups

Parameter	PCOS (n=16)	Non-PCOS (n=44)	p-value
Serum testosterone (ng/dL)	78.5 ± 12.3	42.1 ± 9.8	<0.001*
Fasting insulin (μU/mL)	20.3 ± 4.6	12.5 ± 3.1	<0.001*
Fasting glucose (mg/dL)	94.7 ± 6.1	88.3 ± 5.4	0.01*
HOMA-IR	3.9 ± 1.2	2.1 ± 0.7	<0.001*

The study found a concerning high prevalence of PCOS among adolescent girls in Lahore, with urban girls almost twice as likely to be affected as rural girls. Socioeconomic differences, particularly higher family income and parental education in urban areas, were paradoxically linked with higher obesity rates, greater sedentary behavior, and more frequent fast-food consumption.

The clinical manifestations were dominated by menstrual irregularities and hyperandrogenic signs, including hirsutism and acne. Regardless of their place of residence, PCOS girls had clear evidence of insulin resistance, as they had significantly high testosterone, fasting insulin levels and higher HOMA-IR scores than their non-PCOS peers. In addition, urban girls had higher rates of obesity and metabolic disturbances, which indicate that environmental and lifestyle factors heighten the genetic risk of PCOS in these populations.

DISCUSSION

The prevalence and risk factors of polycystic ovary syndrome (PCOS) in urban and rural adolescent girls of Lahore, Pakistan was studied in this study. In this cohort, PCOS was overall prevalent in 26.7%, and was significantly higher in the urban (36.7%) than the rural (16.7%). This is consistent with other global and regional reports stating that PCOS is a rising public health problem among adolescents, and particularly in South Asia where urbanization and lifestyle changes are transforming health profiles [11, 12].

Several interrelated factors such as higher rates of obesity, sedentary behaviour, and fast-food consumption are responsible for the high prevalence of PCOS in urban girls. In parallel, BMI, waist circumference, and rates of obesity were significantly greater for urban participants in this study than for rural girls, which is consistent with the well-described association between obesity and the pathophysiology of PCOS. While obesity does not appear to worsen hyperandrogenism in PCOS, it does exacerbate insulin resistance to increase the clinical severity of PCOS. In addition, the prevalence of urban girls with higher frequency of sedentary lifestyle and poor dietary patterns increase their metabolic risk [13].

The most common clinical features were menstrual disturbances especially oligomenorrhea which preceded acne, hirsutism and acanthosis nigricans and were more common in PCOS cases in both urban and rural settings [14]. This pattern emphasizes the need to screen adolescent girls with menstrual irregularities for this reason, as early detection can lead to timely intervention and prevent long term complications. Surprisingly, rural girls with PCOS were not metabolically benign, and even in lower resource settings, PCOS manifests a significant health burden [15]. More cases in urban areas reported a positive family history of PCOS, which suggests the influence of complex interaction between genetic predisposition and environmental factors.

The urban environment appears to magnify the expression of genetic risk through

modifiable factors such as diet and activity levels, emphasizing the need for culturally tailored prevention strategies [16].

The study's strengths include its focus on a vulnerable adolescent population, use of the Rotterdam diagnostic criteria, and the inclusion of both urban and rural groups, which allowed meaningful comparison [17]. However, there are limitations. The small sample size, purposive sampling method, and single-city setting limit the generalizability of the findings. Furthermore, the use of transabdominal rather than transvaginal ultrasound may have underestimated ovarian morphology, and self-reported lifestyle data are subject to recall bias [18].

Future research should include larger, multicentre studies using representative sampling to capture the true burden of PCOS in Pakistan. In addition, longitudinal studies are needed to understand the natural history of adolescent PCOS and to evaluate the effectiveness of early lifestyle and pharmacological interventions [19].

CONCLUSION

This study demonstrates a high prevalence of polycystic ovary syndrome among adolescent girls in Lahore, Pakistan, with urban girls disproportionately affected. Obesity, sedentary lifestyle, poor dietary habits, and positive family history emerged as important contributors to the development of PCOS. Early identification, particularly among girls presenting with menstrual irregularities, is critical to prevent long-term metabolic and reproductive complications. Public health strategies focusing on obesity prevention, lifestyle modification, and adolescent health education are urgently needed, particularly in urban areas where the burden appears highest. Addressing PCOS at the adolescent stage offers a unique opportunity to improve women's health trajectories and reduce the future burden

of diabetes, cardiovascular disease, and infertility in Pakistan.

Conflict of Interest:

The authors declare that no conflicts of interest exist.

Funding:

No external funding was received for this study.

Acknowledgments:

We extend our sincere gratitude to our colleagues and paramedical staff for their invaluable support in making this study possible.

Authors' Contributions:

All authors contributed equally to this work.

Data Availability:

De-identified data are available from the corresponding author upon reasonable request.

REFERENCES

1. Sharma P, Mandeep K, and Khetarpal P. CYP19 gene rs2414096 variant and differential genetic risk of polycystic ovary syndrome: a systematic review and meta-analysis. *Gynecological Endocrinology*. 2021;37(2):126-31.doi: 10.1080/09513590.2020.1813274
2. Sharma P, Mandeep K, Sachin K, and Khetarpal P. A cross-sectional study on prevalence of menstrual problems, lifestyle, mental health, and PCOS awareness among rural and urban population of Punjab, India. *Journal of Psychosomatic Obstetrics & Gynecology*. 2022;43(3):349-58.doi: 10.1080/0167482X.2021.1965983
3. Alina R, Um ES, Hafiz Ghulam Murtaza S. Measuring the Awareness of Polycystic Ovarian Syndrome (PCOS) among Women in Punjab, Pakistan. *Scientific Inquiry and Review*. 2023;7(1).doi: 10.32350/sir.71.01
4. Hussain M, Shaikh F, Jamil L, Shaheen A, Afaq MH, Hasni AA, et al. Exploring the Contributing Factors of Polycystic Ovary Syndrome (PCOS). *Indus Journal of Bioscience Research*. 2025;3(1):696-708.doi: 10.70749/ijbr.v3i1.555
5. Sharif M, Kiran Majeed H, Tagar K, Lohana S, Rauf A, Sharif M, et al. Reproductive Health-Related Knowledge, Attitude, and Practices in

- Women of Reproductive Age in Underdeveloped Areas of Punjab, Pakistan. *Cureus*. 2022;14(11):e31043.doi: 10.7759/cureus.31043
6. Taghavi S-A, Aramesh S, Azizi-Kutenaee M, Allan H, Safarzadeh T, Taheri M, et al. The influence of infertility on sexual and marital satisfaction in Iranian women with polycystic ovary syndrome: a case-control study. *Middle East Fertility Society Journal*. 2021;26(1):2.doi: 10.1186/s43043-020-00047-y
 7. Chopra SM, Misra A, Gulati S, Gupta R. Overweight, obesity and related non-communicable diseases in Asian Indian girls and women. *European Journal of Clinical Nutrition*. 2013;67(7):688-96.doi: 10.1038/ejcn.2013.70
 8. Ganie MA, Chowdhury S, Suri V, Joshi BN, Bhattacharya PK, Agarwal S, et al. Variation in normative values of major clinical biochemistry analytes in healthy reproductive-age women in India: A subset of data from a National Indian Council of Medical Research-Polycystic Ovary Syndrome task force study. *Indian Journal of Pharmacology*. 2023;55(2).doi: 10.4103/ijp.ijp_694_22
 9. Dhar S, Mridha S, Bhattacharjee P. Mutational Landscape Screening Through Comprehensive In Silico Analysis for Polycystic Ovarian Syndrome-Related Genes. *Reproductive Sciences*. 2022;29(2):480-96.doi: 10.1007/s43032-021-00752-7
 10. Alshammary AF, Alsobaie SF, Alageel AA, Aldakheel FM, Ansar S, Alrashoudi R, et al. Molecular Role of Asn680Ser and Asp37Glu Missense Variants in Saudi Women with Female Infertility and Polycystic Ovarian Syndrome. *Current Issues in Molecular Biology* [Internet]. 2023; 45(7):[5494-514 pp.].doi: 10.3390/cimb45070348
 11. Shah SM, Mohanty S, Kaduluri CS, Bhatt GS, Ninama P, Bapat NK. A nationwide cross-sectional study to analyse the impact of the COVID-19 pandemic on lifestyle and manifestations of polycystic ovarian syndrome among clinically diagnosed polycystic ovarian syndrome cases aged 15-49 years. *International Journal Of Community Medicine And Public Health*. 2022;10(1):240-9.doi: 10.18203/2394-6040.ijcmph20223551
 12. Ganie MA, Chowdhury S, Malhotra N, Sahay R, Bhattacharya PK, Agrawal S, et al. Prevalence, Phenotypes, and Comorbidities of Polycystic Ovary Syndrome Among Indian Women. *JAMA Network Open*. 2024;7(10):e2440583-e.doi: 10.1001/jamanetworkopen.2024.40583
 13. Shinde KS, Patil SS. Incidence and risk factors of polycystic ovary syndrome among women in reproductive age group attending a tertiary health care hospital in Western Maharashtra. *International Journal of Reproduction, Contraception, Obstetrics and Gynecology*. 2019;8(7):2804-9.doi: 10.18203/2320-1770.ijrcog20193046
 14. Mahar B, Shah T, Sadiq N, Mangi R, Warsi J, Abbas Q. Vitamin D Deficiency Prevalence in Pakistan: Common, Important, and Neglected: A Comprehensive Meta-Analysis. *Journal of Diabetology*. 2024;15(4):335-48.doi: 10.4103/jod.jod_61_24
 15. Balaji S, Amadi C, Prasad S, Bala Kasav J, Upadhyay V, Singh AK, et al. Urban rural comparisons of polycystic ovary syndrome burden among adolescent girls in a hospital setting in India. *Biomed Res Int*. 2015;2015:158951.doi: 10.1155/2015/158951
 16. Rafique A, Salma U, Saleem H. Measuring the Awareness of Polycystic Ovarian Syndrome (PCOS) among Women in Punjab, Pakistan. *Scientific Inquiry and Review*. 2023;7:1-16.doi: 10.32350/sir.71.01
 17. Jena Samanta L, Parida J, Badamali J, Pradhan A, Singh PK, Mishra BK, et al. The incidence, prevalence, and contributing factors of overweight and obesity among adolescent population of India: A scoping review protocol. *PLOS ONE*. 2022;17(9):e0275172.doi: 10.1371/journal.pone.0275172
 18. Zulfiqar S, Noor S, Rafique H, Rehman B, Babar A, Shahid T, et al. Investigation of Prevalence and Awareness of Polycystic Ovary Syndrome among Pakistani Females: Polycystic Ovary Syndrome in Pakistani Women. *Proceedings of the Pakistan Academy of Sciences: B Life and Environmental Sciences*. 2022;59:77-83.doi: 10.53560/PPASB(59-1)703

19. Hayat FA, Shahzad A, Eman U, Malik L, Aziz H, Akbar SUID, et al. Differential Prevalence of Depression Symptoms in Obese Versus Non-obese Women with Polycystic Ovary Syndrome: Investigating the Impact of Obesity on Depression in PCOS Patients. DEVELOPMENTAL MEDICO-LIFE-SCIENCES. 2024;1(3):27-34.doi: 10.69750/dmls.01.03.033

This Article May be cited As: Butt SW, Farooq U, Butt TW, Rai AN, Anjum A. Prevalence and Risk Factors of Polycystic Ovary Syndrome Among Adolescent Girls in Urban and Rural Areas of Punjab, Pakistan: PCOS Risk Factors in Lahore Adolescents. DEVELOPMENTAL MEDICO-LIFE-SCIENCES. 2025;2(4): 21-9.doi: 10.69750/dmls.02.04.0121

Publisher's Note:

Developmental Medico-Life-Sciences remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Developmental Medico-Life-Sciences
Research and Publications Pvt Ltd.**