

Postoperative Pain Management in Patients Undergoing Hernia Repair Surgery. A Cross-Sectional Study

Muhammad Yahya Akhter^{1*}, Muhammad Abdullah Javaid¹, Ayesha Majid¹, Aroosha Ali¹, Sarmad Javed², Eman Tariq¹

1- Lahore Medical & Dental College (LM&DC), Lahore, Pakistan

*Corresponding Author: Muhammad Yahya Akhter, Email: myahya20303@gmail.com, Cell: +923354212348



ABSTRACT

Background: Hernia repair surgery is the most frequently performed surgical operation across the world and pain management after the surgery is important to avoid chronic pain and improve the quality of life.

Objectives: to assess the effectiveness of acetaminophen monotherapy against combination analgesic regimens (acetaminophen plus pethidine or parecoxib) in managing postoperative pain in individuals having hernia repair surgery.

Methods: This was a cross-sectional study on patients who were 18-65 years old and had elective hernia repair surgery. Participants were assigned to three groups: Group A was given IV acetaminophen along with IM pethidine, Group B was given IV acetaminophen with IV parecoxib and Group C was given IV acetaminophen alone. Pain was evaluated by Numeric Rating Scale (NRS) at various time intervals within 24 hours following surgery. Data were analysed employing repeated measures analysis of variance (ANOVA) and multiple regression analysis.

Results: The research had a total of 259 patients. In comparison to Group C, patients in Groups A and B consistently reported far lower pain scores. Group C had the greatest mean pain score (7.91) at 45 minutes post-surgery, whereas Groups A and B demonstrated lower ratings (5.99 and 6.87). Pethidine and parecoxib when paired with acetaminophen were equally effective, as evidenced by the lack of a significant difference between Groups A and B. If acetaminophen monotherapy was used, combination treatment produced better pain relief.

Conclusion: Following hernia repair surgery, combination analgesic regimens comprising acetaminophen and either pethidine or parecoxib are more successful at reducing postoperative pain than acetaminophen monotherapy. As a normal procedure, multimodal analgesia should be taken into account to reduce the incidence of chronic pain and enhance patient recovery.

Keywords: Hernia repair surgery, Postoperative pain management, Acetaminophen, Pethidine, Parecoxib, Multimodal analgesia.



Received: 20/05/2024

Revised: 25/07/2024

Accepted: 29/07/2024

Published: 31/07/2024

© The Author(s) 2024. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third-party material in this article are included in the article's Creative Commons license unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you must obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/public domain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

INTRODUCTION

Hernia repair surgery is one of the most frequent surgeries performed globally, and surgeries are conducted daily, monthly, yearly, and so on. Another major issue following surgery for hernia is pain management because uncontrolled pain after the surgery may result in chronic pain[1]. Chronic postoperative pain has been found to affect 8-16% of the patients who underwent hernia repair surgery and this significantly hampers their overall

functioning[2]. Pain management after surgery is therefore crucial in order to enhance patients' prognosis, reduce hospital stay and prevent chronic pain[3]. Pain management after surgery is greatly dependent on the analgesic regimen that is to be chosen. They include Acetaminophen which has been found to be safer; however, its effectiveness may be quite low when used singly[4]. The co-administration of the non-opioid analgesics with opioids or other analgesics has been

demonstrated to improve the pain relief, decrease the requirement for opioids and, in addition, limit the side effects of opioids[4, 5]. Chronic pain after surgery is a sub-type of chronic pain. It is established that chronic postoperative pain is possible after many types of surgery including amputation, thoracotomy, and inguinal hernia surgery[6]. The development of chronic pain after inguinal hernia repair has been researched; nonetheless, the management is challenging. Groin hernia repair surgeries are performed by millions of times annually around the world. About few percent of these patients suffer from disabling pain and because of the high frequency of groin hernia repairs they present a significant clinical concern[7]. In the case of pain and the extent of the limitation of the activities, the patient should be given an option for an evidence-based treatment plan[8]. One source of criticism which is related to the indices of the current evidence is the variation in the methods of assessment and the length of follow up in the chronic pain after groin hernia repair[9, 10]. Pain management after surgery is an important aspect of patient care, this study aims to assess by comparing the efficacy of combination postoperative analgesic regimens with those of acetaminophen monotherapy in patients with hernia repair surgery[11]. By comparing pain scores and analgesic requirements, this study seeks to provide insights into optimizing postoperative pain management and reducing the incidence of chronic pain.

MATERIALS AND METHODS

This study was undertaken to determine the effectiveness of pain control in surgical patients who underwent hernia repair surgery. This cross-sectional study was carried out on the patients who had undergone open or laparoscopic hernia repair surgery in Ghurki trust & teaching hospital Lahore from September 2022 till May 2023. Pain and measures used in its control were evaluated within 24 hours of the surgical procedure. Patients were stratified according to the American Society of Anesthesiologists (ASA) physical status I or II. Exclusion criteria excluded patients on chronic pain, opioid treatment, or contraindications to the used analgesics. Ethical considerations were followed, written consent was obtained from each participant after a comprehensive description of the study's objective, methods, risks, and potential benefits were given, with ethical issues taken into account. Patients were advised that their participation was completely optional and that doing so would not affect their access to medical care at any point. All patient data were securely maintained and utilized only for research in order to maintain anonymity. In order to ensure adherence to ethical standards and protect participant welfare, the study was granted ethical approval by the institutional review board (IRB) of Lahore University of Biological & Applied Sciences (Lahore-UBAS), a project of Lahore Medical & Dental College (LMDC), Pakistan. Ethical

approval letter ref no. 2023/15C. Patients with a condition of interest adults, between 18 to 65 years of age, involved in the study. Participants who had hernia repair surgery, including inguinal, femoral, umbilical or incisional hernias were included. Both inguinal and femoral and both the open and the laparoscopic repair were considered. Surgical patients who could meet the researcher at the hospital within 24 hours after the surgery and gave a consent to be interviewed about their pain were included. Patients with a history of chronic pain, chronic opioid use, or allergies to NSAIDs or opioids were excluded from the study to avoid confounding variables. The patients were further categorized into three groups in relation to the postoperative analgesic regimen that they received. Group A was given IV acetaminophen 1 g eight hourly and IM pethidine 50 mg six hourly. The members of Group B were given IV acetaminophen 1 g every 8 hours and IV parecoxib 40 mg every 12 hours. The subjects of group C additionally underwent IV acetaminophen monotherapy, in which they were administered 1 g every 8 hours. Patients' pain was measured using the Numeric Rating Scale (NRS) at 45 minutes, 2 hours, 6 hours, 12 hours and 24 hours after surgery. Quantitative data were analyzed using Statistical Package for Social Sciences (SPSS) version 25.0. Repeated-measures ANOVA was used to compare pain scores across groups, with post-hoc tests to identify specific differences. Multiple regression analysis determined the relationship between patient characteristics and pain scores. $p \leq 0.05$ was considered statistically significant.

RESULTS

The demographic and clinical characteristics of the study population are shown in the table 1. The distribution of the patients between the three groups was also balanced and there was no statistical difference of the patients in terms of age, gender and ASA classification. This implies that the groups were balanced and this is important in order to reduce confounding factors that may be present at the time of comparing postoperative pain scores to the different analgesic regimens. The lack of intraoperative complications in all the groups also shows that the surgical procedure was not different in any of the groups, which also enhances the validity of the subsequent pain measurements. The demographic and clinical characteristics of the study population are shown in the (Table- 1).

The mean NRS pain scores depicted in Table 2, it is clear that patients who were given combination therapy of acetaminophen and pethidine or parecoxib reported significantly lower pain scores than those who were given acetaminophen only as in Group C. Ninety minutes after the surgery patients in Group C rated the highest pain level (7.91, on the NRS), while the patients in Group A and Group B had significantly lower pain (5.99 and 6.87, respectively). This increase in pain score was seen

throughout the 24-hour period with Group C having higher scores throughout the period.

Thus, these results give an indication that the combination therapies were more useful in the postoperative pain relief soon after surgery and in the postoperative period. The observed decrease in pain scores of Groups A and B can be explained by the additivity of the used drugs acetaminophen with either an opioid (pethidine) or a non-opioid analgesic (parecoxib) which probably yielded better analgesia comparing with a single agent, acetaminophen.

Table 3 shows pain scores comparison of the groups using statistical analysis. The mean difference of pain scores in Group A (acetaminophen + pethidine) and Group B (acetaminophen + parecoxib) was small (0.12) and P-value was 1.000, therefore showing no significant

difference in the pain-relieving effect between these two combination therapies. However, when comparing mean between group A and C; group B and C, the differences are relatively higher with mean difference of 1.87 and 2.05 respectively and $P < 0.0001$ showed highly significant differences. These results support the findings that both the combination therapies offered better pain relief than just using acetaminophen alone.

Non-significant difference between Groups A and B can be assumed indicating that pethidine has the same efficacy as parecoxib when taken with acetaminophen. This has an implication since it provides a diversity in clinical practice since the healthcare providers can decide on which option to use depending on other factors such as tolerance levels or the contraindications of a certain drug.

Table 1: Demographic and Clinical Characteristics of the Study Population

Characteristic	Group A (Acetaminophen + Pethidine)	Group B (Acetaminophen + Parecoxib)	Group C (Acetaminophen Monotherapy)
Number of Patients (n)	103	94	62
Mean Age (years)	55 (36-65)	52 (35-61)	49 (38-57)
Gender (M/F)	103/30	94/50	62/10
ASA Classification I/II	79/24	72/22	47/15
Mean Operative Time (minutes)	55.2 ± 8.8	59.7 ± 12.6	58.5 ± 9.3

Table 2: Mean NRS Pain Scores at Different Time Points Post-Surgery

Time Point	Group A (Acetaminophen + Pethidine)	Group B (Acetaminophen + Parecoxib)	Group C (Acetaminophen Monotherapy)
45 minutes	5.99 ± 0.75	6.87 ± 0.75	7.91 ± 0.75
2 hours	4.93 ± 0.60	4.66 ± 0.60	6.23 ± 0.60
6 hours	3.54 ± 0.40	3.61 ± 0.40	4.84 ± 0.40
12 hours	2.24 ± 0.30	2.02 ± 0.30	3.89 ± 0.30
24 hours	1.53 ± 0.20	1.39 ± 0.20	2.40 ± 0.20

Table 3: Statistical Analysis of Pain Scores Between Groups

Comparison	Mean Difference	Standard Deviation (SD)	P-Value
Group A vs Group B	0.12	0.05	1.000
Group A vs Group C	1.87	0.10	< 0.0001
Group B vs Group C	2.05	0.10	< 0.0001

DISCUSSION

The findings of this research are in agreement with prior literature that has indicated the advantages of combination analgesic therapy for postoperative pain relief [12]. Multimodal analgesia was shown to be superior to the use of a single agent, because the addition of either pethidine or parecoxib to acetaminophen significantly improved pain scores compared to the administration of acetaminophen alone [13]. This approach is not only beneficial in improving pain and analgesia but also limits the use of opioids diminishing the risk of side effects associated with opioids [14, 15, 16]. The outcomes of the study are significant to the clinical practice in the management of pain after hernia repair surgery. Because chronic pain might develop due to ineffective postoperative analgesia, especially in patients with poor pain management, combination analgesic administration should be considered as best practices [17]. Nevertheless, it is vital to consider

the patient's background, the history of the pain, as well as his/her tolerance levels and possible reaction to the pain medication [18, 19]. It is also worth mentioning, that pain level during mobilization was not evaluated, thus, it might be interesting to see how the analgesic regimens affect the participants during actual physical activity. To the present study, it is recommended that future studies examine pain management during postoperative mobilization and examine the rate of persistent pain after hernia surgery [20]. From the study, it was established that postoperative pain management in hernia repair patients was well managed by the use of NSAIDs in combination with opioids [21].

Patient controlled analgesia was achieved by NSAIDs that offered baseline pain relief but with less inflammation compared to other studies and opioids which were used to address the cases of breakthrough pain, mainly in the open surgical procedures. Patients who underwent lumbar spine surgery who were managed with a

multimodal analgesic plan had better pain control and required less supplementary analgesia[22]. Early and customized analgesic medication delivery proved essential for reducing pain and accelerating healing. Personalized pain management regimens are necessary to improve patient comfort throughout the postoperative period and maximize results, as indicated by the variety in patient reaction[23].

CONCLUSION

The study demonstrated that the combination of analgesic regimens including acetaminophen with pethidine or parecoxib is more effective than the acetaminophen monotherapy in controlling the postoperative pain after hernia repair surgery. Multimodal analgesia should be considered a standard approach to minimize chronic pain incidence and enhance patient recovery.

Funding: None

Conflict of interest: No conflict of interest was declared.

Acknowledgment: We acknowledge the contributions made to this study by all of the team members and participants.

Authors' Contribution: The study was conceptualized by MYA and MAJ. AM and AA gathered the information. ET and SJ analyzed the data. The final version of the paper was approved by all authors after it was reviewed by MYA.

REFERENCES

1. Sinmaz T, Akansel N. Experience of pain and satisfaction with pain management in patients after lumbar disc herniation surgery. *J Perianesth Nurs.* 2021;36(6):647-55. doi:10.1016/j.jopan.2021.01.009
2. Amberbir WD, Bayable SD, Fetene MB. Prevalence and factors associated with acute postoperative pain in elective gynecologic surgical patients in Addis Abeba: a cross-sectional study. *Ann Med Surg.* 2023;85(6):2506-11. doi:10.1097/MS9.0000000000000716
3. Bande D, Moltó L, Pereira JA, Montes A. Chronic pain after groin hernia repair: pain characteristics and impact on quality of life. *BMC Surg.* 2020;20(1):147. doi:10.1186/s12893-020-00805-9
4. Jakobsson E, Lundström KJ, Holmberg H, de la Croix H, Nordin P. Chronic pain after groin hernia surgery in women: patient-reported outcomes from the Swedish Hernia Register. *Ann Surg.* 2022;275(2):213-9. doi:10.1097/SLA.0000000000005194
5. Nishikawa M, Fukuda T, Okazaki M. Predictive factors of postoperative acute pain in laparoscopic inguinal hernia repair in men: a single-centre retrospective study. *J Perioper Pract.* 2023;33(5):133-8. doi:10.1177/17504589211054371
6. Singh PK, Saikia P, Lahakar M. Prevalence of acute postoperative pain in adults undergoing abdominal surgery and correlation with analgesic satisfaction. *Indian J Anaesth.* 2016;60(10):737-43. doi:10.4103/0019-5049.191686
7. Özdemir C, Karazeybek E, Söyüncü Y. Relationship between care quality and postoperative pain outcomes in major orthopedic surgery. *Clin Nurs Res.* 2022;31(3):530-40. doi:10.1177/10547738211059960
8. Deng X, Liang S, Li H, Gouda D, Zhu T, Xiao K. Perception differences of day-surgery information among patients and staff. *Patient Prefer Adherence.* 2019;13:381-7. doi:10.2147/PPA.S196674
9. Olsson A, Sandblom G, Fränneby U, Sondén A, Gunnarsson U, Dahlstrand U. Postoperative complications and risk of chronic groin pain after open inguinal hernia repair. *Surgery.* 2017;161(2):509-16. doi:10.1016/j.surg.2016.08.011
10. Smith L, Meggy A, Watts T, Knight L, Torkington J, Cornish J. Incisional hernia prevention: INVITE protocol. *BMJ Open.* 2022;12(12):e069568. doi:10.1136/bmjopen-2022-069568
11. Malibary N, Shurrab M, Albariqi MO, Bohairi M, Basabain AS, Alqurashi MY, et al. Quality of life after umbilical hernia repair. *Cureus.* 2021;13(10):e19016. doi:10.7759/cureus.19016
12. López-Cano M, Rodrigues-Gonçalves V, Verdaguier-Tremolosa M, et al. Prioritization criteria for abdominal wall hernia surgery. *Hernia.* 2021;25(6):1659-66. doi:10.1007/s10029-021-02378-9
13. Ndong A, Diallo AC, Diao ML, et al. Acute postoperative complications increase recurrence and chronic pain after inguinal hernia surgery. *Int J Abdom Wall Hernia Surg.* 2023;6(4):236-41. doi:10.4103/ijawhs.ijawhs_47_23
14. Bragais LCG, Faylona JMV. Adherence to groin hernia management guidelines: cross-sectional study. *Hernia.* 2020;24(5):969-75. doi:10.1007/s10029-020-02207-5
15. Zeleke TG, Mersha AT, Endalew NS, Ferede YA. Back pain after spinal anesthesia: cross-sectional study. *Adv Med.* 2021;2021:6654321. doi:10.1155/2021/6654321
16. Liu Y, Xiao S, Yang H, et al. Postoperative pain outcomes and perioperative pain management in China. *Lancet Reg Health West Pac.* 2023;39:100822. doi:10.1016/j.lanwpc.2023.100822
17. Menlah A, Osei-Appiah E, Garti I, Frempomaa Agyare D. Factors influencing postoperative pain management among nurses. *J Perioper Pract.* 2023;33(7-8):223-32. doi:10.1177/17504589211064039
18. Demelash G, Berhe YW, Gebregzi AH, Chekol WB. Post-cesarean postoperative pain: observational study. *Open Access Surg.* 2022;15:1-8. doi:10.2147/OAS.S347920
19. Slooter CD, Perquin CW, Zwaans WA, et al. Laparoscopic mesh removal for chronic post-hernia pain. *Hernia.* 2023;27(1):77-84. doi:10.1007/s10029-022-02712-9
20. Torensma B, Hany M, Bakker MJS, et al. Pre- and postoperative chronic pain in bariatric surgery: e-survey. *Obes Surg.* 2023;33(1):204-10. doi:10.1007/s11695-022-06354-9
21. Bjurström MF, Irwin MR, Chen DC, Smith MT, Montgomery A. Sex differences, sleep disturbance, and persistent pain after groin hernia surgery. *J Pain.* 2021;22(11):1360-70. doi:10.1016/j.jpain.2021.04.008
22. Chu Z, Zheng B, Yan L. Chronic pain after inguinal hernia surgery: systematic review and meta-analysis. *Hernia.* 2024;28(4):967-87. doi:10.1007/s10029-024-02980-7
23. Bozkurt H, Gür HÜ, Akıncı M, Aslan H, Karakullukçu Ç, Yıldırım D. Emergency surgery in a COVID-19 hospital: cross-sectional study. *Sao Paulo Med J.* 2020;138.

This Article May be cited As: Akhter MY, Javaid MA, Majid A, Ali A, Javed S, Tariq E. Postoperative Pain Management in Patients Undergoing Hernia Repair Surgery: A Cross-Sectional Study. Dev Med Life Sci. 2024;1(5):20-25. doi: 10.69750/dmls.01.05.052.

Publisher's Note:

Developmental Medico-Life-Sciences remains neutral with regard to jurisdictional claims in published maps. and institutional affiliations.

