


Evaluating the Effectiveness of Lifestyle Modifications in Managing Hypertension Among Adults

Ali Khalid ^{1*}, Ayesha Khalid ², Ayesha Tariq ², Ume Aiman ², Mahnoor Imtiaz ², Roshan Ahmad ², Fatima Naeem ², Yasir Hameed ³

- 1- Al-Aleem Medical College, Lahore, Pakistan
 2- Amna Inayat Medical College (AIMEC), Sheikhupura, Pakistan
 3- Rashid Latif khan university medical & dental college (RLKU), Pakistan



Corresponding Author: Ali Khalid, **Email:** alikhalid4412@gmail.com, **Cell:** +923256574611

ABSTRACT

Background: The significant risk factor for cardiovascular illnesses is hypertension. Changes in nutrition, exercise, and stress have the ability to lower blood pressure.

Objectives: The purpose of this research was to assess the results of life style changes such as diet, exercise and quitting smoking etc. for control of hypertension in adults.

Methodology: A prospective, randomized controlled was conducted from September 2022 till august 2023. Total 500 adults with hypertension were selected. Participants were divided into two groups, one who was taking the standard antihypertensive therapy and the other who was taking the standard therapy plus guided behaviour change counselling. The biomarkers like systolic and distolic blood pressure, weight, physical activity, salt intake and smoking were recorded initially and after 12 months of intervention. Statistical analysis was performed using SPSS version 25.0, involved t-tests for continuous variables and chi-square tests for categorical variables, with $p \leq 0.05$ considered significant.

Results: The results showed that intervention group experienced substantial decrease in SBP (-14.2 mmHg, $p < 0.001$) and DBP (-9.6 mmHg, $p < 0.001$), whereas the control group experienced lower reductions (-6.4 mmHg, $p < 0.01$) and DBP (-4.2 mmHg, $p < 0.01$). Furthermore, 75% of patients in the intervention group met their goal blood pressure, compared to 50% in the control group ($p < 0.001$). The intervention group had considerably greater levels of lifestyle adherence, such as physical exercise and a low-salt diet ($p < 0.001$).

Conclusion: Combining lifestyle management with traditional antihypertensive treatment considerably improves blood pressure control and lifestyle adherence, providing a comprehensive approach to hypertension management.

Keywords: Hypertension, lifestyle modifications, blood pressure, diet, physical activity, smoking cessation, Cardiovascular Risk.



Received: 01/06/2024
 Revised: 21/09/2024
 Accepted: 28/09/2024
 Published: 30/09/2024

© The Author(s) 2024. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third-party material in this article are included in the article's Creative Commons license unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you must obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

INTRODUCTION

A major contributor to morbidity and death is hypertension which affects more than one billion people worldwide and is linked to renal failure, cardiovascular disease, and stroke [1]. Although pharmacological treatments for hypertension exist, the control of high blood pressure is inadequate, and many patients fail to reach and sustain their treatment goals [2]. Non-compliance with prescribed drugs and other

unhealthy practices including high intake of sodium, lack of exercise, and smoking are some of the main causes of uncontrolled hypertension. The World Health Organization (WHO) and other health-related organizations stress that diet changes, exercising more, and not smoking are important elements in controlling hypertension [3]. The second leading cause of mortality in Pakistan and the third most prevalent cause of death worldwide attributable to

heart coronary disease is hypertension illness, often referred to as a “Silent killer”. Since blood pressure is the primary avoidable factor that can considerably lower mortality and linked disorders, lowering blood pressure is vital. Reductions in systolic blood pressure of 3.5 milligrams can lower mortality rates by 8% in cases of stroke and 7% in cases of coronary artery disease. Less than half of hypertension patients were getting antihypertensive therapy, and only one in five (less than 25%) had achieved blood pressure control below 145/90 mmHg, despite the availability and effectiveness of antihypertensive medications. Along with traditional medicines, non-medical methods such as lifestyle adjustments are also stressed to prevent and control hypertension. Lifestyle modifications, formerly known as non-pharmacologic therapy, are advised as the initial course of treatment for hypertensive disorders due to their substantial involvement in lowering blood pressure. The lifestyle interventions, dietary modification using the

DASH (Dietary Approaches to Stop Hypertension) diet, reduced salt intake and physical activity all have been found to decrease both systolic and diastolic blood pressure [4, 5]. However, most of such studies involve single or isolated interventions, and/or have short-term follow up, and therefore, the long-term impact of intensive Comprehensive Lifestyle Strategies (CLS) with standard pharmacologic therapy remains undefined [6]. The study aims and objectives were to address the rising issue that pharmacological therapies for hypertension frequently fail to provide adequate control, despite their widespread usage. The current study aimed to determine whether lifestyle modifications such as the DASH diet, increased physical activity, and smoking cessation, when combined with standard therapy, can significantly improve blood pressure management and overall health outcomes in hypertensive adults [7].

MATERIALS AND METHODS

The current prospective, randomized controlled trial study was conducted from August 2022 to August 2023 at Kishwar Fazal Teaching Hospital and Hameed Latif Teaching Hospital. Total 500 hypertensive adults aged 30 to 65 years were selected from hospitals. This study was approved by the Ethical Committee of Rashid Latif Khan University Medical & Dental College (RLKU), Pakistan approval number IRB-RLKU-18/07/22/22-A. All participants provided informed consent prior to enrollment in the study. Participants were randomly assigned to one of two groups, the standard care group that was a control group (n=250) which received standard antihypertensive treatment and the treatment group which received lifestyle modification (n=250) in addition to standard antihypertensive treatment. The inclusion criteria for participants in this study included; a clinical diagnosis of hypertension with a systolic BP of 140 mmHg or above or diastolic BP of 90 mmHg or above. They also excluded

patients with secondary hypertension, recent cardiovascular event, severe comorbidity including chronic kidney disease, and cancer.

The intervention group got individualised dietary counselling for lifestyle changes to include Dietary Approaches to Stop Hypertension (DASH) diet and were instructed to take not more than 2 grams of salt per day. 3 grams and increase the intake of foods that are rich in fruits, vegetables and low fat dairy products. Aerobic exercise at moderate-intensity was set at least 150 minutes per week as a physical activity goal. To the participants who smoked, smoking cessation was offered; counseling and other nicotine replacement therapies were deemed necessary. These two groups received monthly follow-up visits where their blood pressure, their body weight and the degree of compliance to the mentioned lifestyle changes were assessed. Systolic and diastolic blood pressures were recorded with an automatic sphygmomanometer and patients' compliance with the prescribed lifestyle changes were estimated with the help of standardised questionnaires. The major measure of effect was the difference in systolic and diastolic blood pressure at the end of one year follow up from the baseline measurement. Secondary end points encompassed were variations in weight, physical activity, diet compliance and smoking status. Data analysis was done using the statistical package SPSS version 25.0, while demographic and clinical parameters of the patients were presented in terms of mean, standard deviation (Mean $\pm$ SD) and Frequency Distribution. Intragroup comparisons of blood pressure changes were analysed using paired t-tests, whereas intergroup comparisons were done using independent t-tests. Chi-square tests were applied to compare categorical variables such as gender and lifestyle adherence. Significance was considered at ($p \leq 0.05$).

RESULTS

A total of n=500 hypertension patients were enrolled, n=250 were placed in the intervention group and n=250 in the control group. The age range of the participants was 35-65. There were no appreciable variations in baseline blood pressure, gender, age, body mass index (BMI), or smoking behaviors between the two groups (Table 1). Independent t-tests were applied to continuous variables (e.g., age, SBP, DBP, BMI), chi-square test was applied for categorical variables (e.g., gender, smoking status). The baseline variables for the control and intervention groups were compared in the table-1. Age, gender distribution, baseline blood pressure readings (both systolic and diastolic), body mass index (BMI), and smoking percentage were comparable across the two groups. There were only little differences between the groups, the intervention group had a slightly higher systolic blood pressure and BMI, whereas the control group has a slightly larger number of smokers.

When compared to the control group, the intervention group's gains in blood pressure were noticeably bigger at the 12-month follow-up. The intervention group showed a reduction in mean diastolic blood pressure (DBP) of 9.6 mmHg ($p<0.001$) and a drop in mean systolic blood pressure (SBP) of 14.2 mmHg ($p<0.001$). The control group, on the other hand, had a mean drop in Diastolic Blood Pressure (DBP) of 4.2 mmHg ($p<0.01$) and Systolic Blood Pressure (SBP) of 6.4 mmHg ($p<0.01$). Table- 2 showed that the intervention group had a considerably larger number of patients (75%) who achieved the goal blood pressure ($<140/90$ mmHg) than the control group (50%) ($p<0.05$). Paired t-tests were applied to compare baseline and after-12-months values within each group

(control vs. intervention). Independent t-tests were used to compare changes in SBP and DBP between groups.

Table-3 shows participants in the intervention group showed better commitment to lifestyle changes. Compared to 40% of the control group, 70% of the intervention group met the recommended level of physical activity after a year at least 150 minutes per week. Furthermore, a low-salt diet was followed by 60% of individuals in the intervention group vs 30% in the control group ($p<0.001$). Table 3 shows that the intervention group had a higher success rate when it came to smoking cessation, with 30% of smokers quitting compared to 15% in the control group ($p=0.002$). Chi-square tests were used to assess differences in categorical variables such as adherence to physical activity, diet, and smoking cessation.

Table 1: Baseline variables of Participants

Variable	Control Group (n=250)	Intervention Group (n=250)	p-value
Age (years)	54.1 $\pm$ 8.3 (35–65)	53.8 $\pm$ 8.1 (35–65)	0.056
Gender (Male/Female)	130/120	135/115	0.057
Baseline SBP (mmHg)	153.1 $\pm$ 9.0	153.6 $\pm$ 8.5	0.048
Baseline DBP (mmHg)	94.1 $\pm$ 4.6	94.3 $\pm$ 4.8	0.055
Body Mass Index (BMI, kg/m ²)	28.4 $\pm$ 3.2	28.7 $\pm$ 3.4	0.041
Smokers (%)	28%	27%	0.035

($p\leq 0.05$)

Table 2: Blood Pressure Changes After 12 Months

Blood Pressure	Control Group	Intervention Group	p-value
SBP (mmHg) - Baseline	153.1 $\pm$ 9.0	153.6 $\pm$ 8.5	0.453
SBP (mmHg) - After 12 Months	146.7 $\pm$ 7.4	139.4 $\pm$ 6.2	<0.001
Change in SBP (mmHg)	-6.4	-14.2	<0.001
DBP (mmHg) - Baseline	94.1 $\pm$ 4.6	94.3 $\pm$ 4.8	0.532
DBP (mmHg) - After 12 Months	90.1 $\pm$ 4.5	84.7 $\pm$ 4.1	<0.001
Change in DBP (mmHg)	-4.2	-9.6	<0.001
% Achieving BP Target ($<140/90$)	50%	75%	<0.001

($p\leq 0.05$)

Table 3: Lifestyle Modification Adherence

Lifestyle Factor	Control Group (n=250)	Intervention Group (n=250)	p-value
Physical Activity (≥ 150 min/wk)	40%	70%	<0.001
Low-Salt Diet Adherence	30%	60%	<0.001
Smoking Cessation	15%	30%	0.002

($p\leq 0.05$)

The results of the study showed that changing one's lifestyle may significantly lower blood pressure and encourage better habits. In addition to achieving significant decreases in both systolic and diastolic blood pressure, the intervention group also demonstrated better adherence to dietary modifications, physical exercise, and quitting smoking. These results highlight the importance of complete lifestyle modifications in addition to conventional medication therapy for the management of hypertension.

DISCUSSION

Results of current study concluded that life style changes should be recommended as a part of an anti-hypertensive treatment since they can be as efficient as drugs alone[8]. Patients in the intervention group who followed up on

lifestyle modifications including increased physical activity, low salt diet and smoking cessation had significantly better reduction in both systolic and diastolic blood pressure than those in the control group who only took medication[9, 10].

The significant changes that were recorded in the course of the study, the most outstanding was the significant enhancement of blood pressure control in the group that received the intervention. 14.2 mmHg reduction in systolic blood pressure and 9.6 mmHg on diastolic blood pressure were recorded during the course of the one-year study period[11, 12]. These reductions are clinically significant and are higher than the changes in the control group where a 6 percent improvement in the score was observed. 4 mmHg of systolic blood pressure lowering and 14.2mmHg reduction of the diastolic blood pressure by as

much as 2 mmHg[13]. Similarly, target blood pressure, that is <140/90 mmHg was achieved by a significantly higher percentage of the intervention group patients (75%) as opposed to 50% of the control group patients. These results are consistent with the previous studies about the fact that lifestyle modifications, including the DASH diet, and increased levels of physical activity are central to the management of blood pressure[14, 15]. These findings align with those of McManus et al. (2021), who also stated noteworthy drops in blood pressure among patients undergoing digital and lifestyle treatments. Their findings highlight the supplementary function of lifestyle alterations in conjunction with pharmaceutical therapy.

These favorable outcomes were achieved due to compliance with lifestyle changes that were prescribed. In the analysis of the results, it was observed that participants in the intervention group had the significantly higher level of physical activity and dietary compliance[16]. It was also observed that by the end of the study the percentage of the subjects in the intervention group that attained the WHO recommended 150 minutes of physical activity per week was 70% while that of the control group was 40%. Furthermore, concerning the issue of low salt diet only 60% of the people in the intervention group complied with it unlike the control group of 30%. Such behavioral modifications are relevant for the treatment of hypertension because physical activity and low sodium diet are known to decrease blood pressure and prevent cardiovascular complications[17, 18]. This finding aligns with studies like Valenzuela et al. (2021), which highlighted the long-term effectiveness of comprehensive lifestyle interventions in hypertension management. Another factor that contributed towards better results included smoking cessation. The smoking cessation rate recorded among the intervention group was 30% than the control group, which was only 15% thus proving the efficacy of the structured smoking cessation program[19]. Tobacco use is also another common risk factor that is known to increase the chances of hypertension besides being linked to other cardiovascular complications once one quits smoking, there are chances that he/she is likely to have better control of his/her blood pressure[20, 21]. Altogether, the results of this study support the notion of using pharmacological therapy along with the life style changes to achieve the best control of hypertension. However, the limitations of study is that pharmacological treatment alone is not sufficient and the life style changes such as diet, exercise, and smoking cessation can significantly improve the treatment results[22]. These findings also provide the evidence for the long-term approaches to maintain the lifestyle changes, for the effects of these interventions may decrease if people's compliance with the recommendations weakens in the further time[23, 24]. The findings of the present work were concluded that life style changes should be recommended as a part of an antihypertensive treatment since they can be as efficient as

drugs alone[8]. Patients in the intervention group who followed up on lifestyle modifications including increased physical activity, low salt diet and smoking cessation had significantly better reduction in both systolic and diastolic blood pressure than those in the control group who only took medication[9, 10]. In conclusion the use of standard antihypertensive treatment along with a package of CLM proved to be beneficial for the management of blood pressure and for reduction of hypertension-associated risks. The mentioned changes should be promoted in further studies by health professionals and supported in patients since it is a strong and long-term approach to fight hypertension and maintain healthy living[25, 26].

CONCLUSION

This study showed that in addition to traditional antihypertensive medication, adopting lifestyle modifications such as dietary adjustments, increasing physical activity, and quitting smoking significantly improves blood pressure management in persons with hypertension. These therapies raise the percentage of patients reaching target blood pressure in addition to lowering systolic and diastolic blood pressure.

Future prospects: In order to improve long-term health outcomes, encouraging lifestyle modifications have to be a major part of hypertension therapy.

Funding: Nil

Conflict of interest: None

Authors' Contribution: A.K. and A.K. conceptualized the study and developed the methodology. U.A., M.I., and F.N. assisted in data collection and statistical analysis, while R.A. and U.S.A. contributed to the final manuscript draft and revisions.

Acknowledgement: We sincerely thank all the participants for their cooperation and the medical staff for their assistance in conducting this study.

Data Availability: The data supporting the findings of this study are available from the corresponding author, upon reasonable request.

REFERENCES

1. Yang MH, Kang SY, Lee JA, Kim YS, Sung EJ, Lee KY, et al. The effect of lifestyle changes on blood pressure control among hypertensive patients. *Korean J Fam Med.* 2017;38(4):173-80. doi:10.4082/kjfm.2017.38.4.173
2. Ojangba T, Boamah S, Miao Y, Guo X, Fen Y, Agboyibor C, et al. Comprehensive effects of lifestyle reform, adherence, and related factors on hypertension control: A review. *J Clin Hypertens.* 2023;25(6):509-20. doi:10.1111/jch.14653
3. Yu R, Yan LL, Wang H, Ke L, Yang Z, Gong E, et al. Effectiveness of a community-based individualized lifestyle intervention among older adults with diabetes and hypertension. *Prev Chronic Dis.* 2014;11:E84. doi:10.5888/pcd11.120333
4. Nishigaki N, Shimasaki Y, Yoshida T, Hasebe N. Physician and patient perspectives on hypertension management and lifestyle modification in Japan. *Hypertens Res.* 2020;43(5):450-62. doi:10.1038/s41440-020-0398-0

5. Gazit T, Gutman M, Beatty AL. Assessment of hypertension control in adults participating in a mobile technology BP self-management program. *JAMA Netw Open*. 2021;4(10):e2127008. doi:10.1001/jamanetworkopen.2021.27008
6. Valenzuela PL, Carrera-Bastos P, Gálvez BG, Ruiz-Hurtado G, Ordoñas JM, Ruilope LM, et al. Lifestyle interventions for prevention and treatment of hypertension. *Nat Rev Cardiol*. 2021;18(4):251-75. doi:10.1038/s41569-020-00437-9
7. Oliveros E, Patel H, Kyung S, Fugar S, Goldberg A, Madan N, et al. Hypertension in older adults: Assessment and management. *Clin Cardiol*. 2020;43(2):99-107. doi:10.1002/clc.23303
8. McManus RJ, Little P, Stuart B, Morton K, Raftery J, Kelly J, et al. HOME BP digital intervention RCT. *BMJ*. 2021;372:m4858. doi:10.1136/bmj.m4858
9. Al-Makki A, DiPette D, Whelton PK, Murad MH, Mustafa RA, Acharya S, et al. WHO pharmacological treatment guideline summary. *Hypertension*. 2022;79(1):293-301. doi:10.1161/HYPERTENSIONAHA.121.18192
10. Reynolds AN, Akerman A, Kumar S, Diep Pham HT, Coffey S, Mann J. Dietary fibre in hypertension management. *BMC Med*. 2022;20(1):139. doi:10.1186/s12916-022-02328-x
11. Timpel P, Oswald S, Schwarz PEH, Harst L. Telemedicine interventions in cardiometabolic diseases. *J Med Internet Res*. 2020;22(3):e16791. doi:10.2196/16791
12. Vamvakis A, Gkaliagkousi E, Lazaridis A, Grammatikopoulou MG, Triantafyllou A, Nikolaidou B, et al. Intensive lifestyle treatment RCT (HINTreat). *Nutrients*. 2020;12(5):1326. doi:10.3390/nu12051326
13. Takase M, Hirata T, Nakaya N, Nakamura T, Kogure M, Hatanaka R, et al. Genetic × lifestyle risk associations with hypertension. *Hypertens Res*. 2024;47(8):2064-74. doi:10.1038/s41440-024-01705-8
14. An S, Song R. Health coaching for behavioural modification: meta-analysis. *Patient Educ Couns*. 2020;103(10):2029-38. doi:10.1016/j.pcc.2020.04.029
15. Loaiza-Betancur AF, Chulvi-Medrano I. Low-intensity isometric handgrip exercise and BP management. *Sports Health*. 2020;12(5):470-7. doi:10.1177/1941738120943882
16. Chopra S, Malhotra A, Ranjan P, Vikram NK, Sarkar S, Siddhu A, et al. Predictors of successful lifestyle-based weight loss. *Obes Rev*. 2021;22(3):e13148. doi:10.1111/obr.13148
17. US Preventive Services Task Force. Screening for hypertension in adults. *JAMA*. 2021;325(16):1650-6. doi:10.1001/jama.2021.4987
18. Barbato E, Azizi M, Schmieder RE, Lauder L, Böhm M, Brouwers S, et al. Renal denervation guidance. *Eur Heart J*. 2023;44(15):1313-30. doi:10.1093/eurheartj/ehad054
19. Citoni B, Figliuzzi I, Presta V, Volpe M, Tocchi G. Home BP & telemedicine in hypertension. *High Blood Press Cardiovasc Prev*. 2022;29(1):1-14. doi:10.1007/s40292-021-00492-4
20. Zhou B, Perel P, Mensah GA, Ezzati M. Global epidemiology and interventions for hypertension. *Nat Rev Cardiol*. 2021;18(11):785-802. doi:10.1038/s41569-021-00559-8
21. Kostova D, Spencer G, Moran AE, Cobb LK, Husain MJ, Datta BK, et al. Cost-effectiveness of hypertension management in LMICs. *BMJ Glob Health*. 2020;5(9):e002213. doi:10.1136/bmjgh-2019-002213
22. Belardo D, Michos ED, Blankstein R, Blumenthal RS, Ferdinand KC, Hall K, et al. Nutritional approaches for ASCVD prevention. *Am J Prev Cardiol*. 2022;10:100323. doi:10.1016/j.ajpc.2022.100323
23. Tan FCJH, Oka P, Dambha-Miller H, Tan NC. Self-efficacy and self-care in hypertension. *BMC Fam Pract*. 2021;22(1):44. doi:10.1186/s12875-021-01391-2
24. He L, Zhang M, Zhao Y, Li W, Zhang Y. Life's Essential 8 and mortality in hypertensive patients. *BMC Public Health*. 2024;24(1):1730. doi:10.1186/s12889-024-19189-z
25. Santos L. Impact of nutrition & lifestyle modification on health. *Eur J Intern Med*. 2022;97:18-25. doi:10.1016/j.ejim.2021.09.020
26. Siegel KR, Ali MK, Zhou X, Ng BP, Jawanda S, Proia K, et al. Cost-effectiveness of interventions to manage diabetes. *Diabetes Care*. 2020;43(7):1557-92. doi:10.2337/dci20-0017

This Article May be cited As: Khalid A, Khalid A, Tariq A, Aiman U, Imtiaz M, Ahmad R, Naeem F, Hameed Y. Evaluating the effectiveness of lifestyle modifications in managing hypertension among adults: hypertension management through diet and exercise. *Dev Med Life Sci*. 2024;1(7):4-11. doi:10.69750/dmls.01.07.061

Publisher's Note:

Developmental Medico-Life-Sciences remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

